

Optical and X-ray Spectral Properties of the Swift BAT-detected AGN

Lisa M. Winter^{1,2}

Richard Mushotzky³, Karen Lewis⁴, Sylvain Veilleux³,
Mike Koss³, Brian Keeney²

¹Hubble Fellow


²CASA/University of Colorado, Boulder, CO

³University of Maryland, College Park, MD

⁴Dickinson College, Carlisle, PA


Swift 2009

9-month BAT Survey



- 153 AGN detected at $\geq 4.8\sigma$
- $F_{14-195\text{keV}} > \text{few} \times 10^{-11} \text{ erg s}^{-1} \text{ cm}^{-2}$
- $< z > \approx 0.03$
- See Tueller et al. (2008)
- The Survey is continuing, with the completed 22 month (262 sources, ApJS, accepted) and on-going 36 month catalogs.
- Sensitivity is still scaling as \sqrt{t}


X-ray Sources



- Collected X-ray properties from Swift XRT, ASCA, *XMM-Newton*, and *Chandra*
- Sources selected:
 - Detected with BAT as a 5σ or higher detection
 - Optical counterpart clearly seen in DSS/2MASS images
- See Winter et al. (2009)

(Left) XMM-Newton contours on DSS images.

Color-Color Plot





Blue = $\log n_H \geq 23$
Green = $23 > \log n_H > 22$
Red = $\log n_H < 22$
Black = Blazar/BL Lac

Full spectral analysis in
Winter et al. (2009).





Among Our Results ...







- Higher fraction of absorbed sources at low luminosity/accretion rate
- The average X-ray spectrum (0.6–10 keV) replicates the CXB slope of 1.4
- Very few Compton thick sources



Optical Study



- Covered 80% of the ‘Northern’ BAT sources
- Spectra from archived SDSS observations (27), our own KPNO 2.1-m observations (40), and the literature (5)
- Half of the spectra shown broad H-Balmer lines





Distribution of Optical Reddening





- Broad line sources: $E(B-V) = 0.08 \pm 0.11$
- Narrow line sources: $E(B-V) = 0.29 \pm 0.33$

Reddening Estimates





- No correlation between optical reddening and host inclination.
- No correlation between optical and X-ray extinction for narrow line sources.
- Unlike the results of Alonso-Herrero et al. (1997), most Sy 1s have more extinction in the optical than X-ray band.

Temperature/Density Diagnostics





- Same distributions of the ratio of [S II] $\lambda 6716/\lambda 6731$, which indicates density
- $N_e \approx 10^4 \text{ cm}^{-3}$, consistent with Koski (1978)
- Different distributions of the ratio of [O III] $(\lambda 4959 + \lambda 5007)/\lambda 4363$, which indicates temperature
- Different temperatures or different densities probed by [O III] $\lambda 4363$?

Comparisons of Mass Determination Methods



- Reverberation mapping/H- β derived masses are well-correlated.
- 2MASS-derived/H- β derived masses are also well-correlated.
- Average mass of the Swift BAT-detected AGN: $\langle M/M_\odot \rangle = 10^{7.87 \pm 0.66}$, consistent with previous studies (Woo & Urry, 2002)

Luminosity Distributions



Broad Line: $\log L_{\text{14--195keV}} = 43.74$, $\log L_{[\text{OIII}]} = 41.79$



Narrow Line: Seyferts: $\log L_{\text{14--195keV}} = 43.87$, $\log L_{[\text{OIII}]} = 41.55$

LINERs: $\log L_{\text{14--195keV}} = 43.50$, $\log L_{[\text{OIII}]} = 40.73$

Others: $\log L_{\text{14--195keV}} = 42.69$, $\log L_{[\text{OIII}]} = 40.33$





X-ray/Optical Luminosity Comparison



- Slight correlations ($R^2 < 0.4$) are seen between [O III] and BAT luminosities: $L_{[OIII]}(\text{corr}) \propto L_{\text{BAT}}^{1.16 \pm 0.24}$. There is a lot of scatter.
- Agrees with Meléndez et al. (2008), in contrast to Heckman et al.




Soft X-ray and Optical Luminosity



- Best correlation ($R^2 = 0.6$) between the [O III] and soft X-ray flux for narrow line sources
- Same seen in the XMM sample presented by Terashima (2009)



Host Galaxy Properties



Narrow Line Sources
Broad Line Sources

- $D_n(4000)$: Old stars through the Ca II break
- $H\delta_A$: Young stars through $H\delta$ absorption
- Narrow Line sources are consistent with intermediate/old populations
- Broad Line sources have lower H δ EWs
- Low EWs (< 0) associated with very young populations of < 0.1 Gyr (Leitherer et al., 1999)

Summary

- We have completed analyses of the X-ray (0.3–10 keV) and optical spectra of the Swift BAT-detected AGN in the 9-month catalog.
- Optical extinction is not from the host galaxy and is correlated with X-ray extinction for Sy 1s but not Sy 2s.
- Optically identified Seyferts have the same distributions of both $[\text{O III}] \lambda 5007\text{\AA}$ and 14–195 keV luminosities for narrow and broad line sources. **This is in agreement with the Unified Model.**
- Correlations between $[\text{O III}] \lambda 5007\text{\AA}$ and 14–195 keV luminosities are weak with much scatter. **$L_{[\text{OIII}]}$ is not the best indicator of L_{bol} .**
- Broad Line sources appear to have much younger stellar host populations than narrow line sources (based on Lick indices).
This conflicts with the Unified Model?

References

- Alonso-Herrero, A., Ward, M. J., & Kotilainen, J. K. 1997, *MNRAS*, 288, 977
- Crenshaw, D. M., Kraemer, S. B., Boggess, A., Maran, S. P., Mushotzky, R. F., & Wu, C.-C. 1999, *ApJ*, 516, 750
- Dunn, J. P., Crenshaw, D. M., Kraemer, S. B., & Gabel, J. R. 2007, *AJ*, 134, 1061
- George, I. M., Turner, T. J., Netzer, H., Nandra, K., Mushotzky, R. F., & Yaqoob, T. 1998, *ApJS*, 114, 73
- Heckman, T. M., Ptak, A., Hornschemeier, A., & Kauffmann, G. 2005, *ApJ*, 634, 161
- Koski, A. T. 1978, *ApJ*, 223, 56
- Leitherer, C. et al. 1999, *ApJS*, 123, 3
- Meléndez, M. et al. 2008, *ApJ*, 682, 94
- Reynolds, C. S. 1997, *MNRAS*, 286, 513
- Tueller, J., Mushotzky, R. F., Barthelmy, S., Cannizzo, J. K., Gehrels, N., Markwardt, C. B., Skinner, G. K., & Winter, L. M. 2008, *ApJ*, 681,

